

Special Populations

Frequent Adverse Effects

Infrequent: Abnormality of accommodation, conjunctivitis, dry eyes, ear pain, photophobia, taste perversion, tinnitus.
Rare: Dizziness, lacrimation disorder, oscillopsia, parosmia, ptosis, strabismus, taste loss, vertigo, visual field defect.

Urogenital System

Infrequent: Abnormal ejaculation, hematuria, impotence, menorrhagia, polyuria, urinary incontinence.
Rare: Acute kidney failure, anorgasmia, breast abscess, breast neoplasm, creatinine increase, cystitis, dysuria, epididymitis, female ejaculation, kidney failure, kidney pain, nocturia, urinary retention, urinary urgency.

5.3 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of SURVEINTE. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Blood and Lymphatic

Aggravated/occasional: hemolytic anemia, lymphadenopathy not associated with hypersensitivity disorder, pseudomyeloma.

Gastrointestinal

Esophagitis

Hypoglycemia/Tyrosinemia and Pancreas

Pancreatitis

Immunologic

Hypomagnesaemia: lupus-like reaction, vasculitis.

Lower Respiratory

Arterio

Musculoskeletal

Arthralgia/myalgia: has been observed in patients experiencing hypersensitivity reactions.

Nervous System

Exacerbation, exacerbation of Parkinsonism symptoms in patients with pre-existing Parkinson's disease, tics.

Non-site Specific

Progressive immunosuppression

Renal and Urinary Disorders

Tubulointerstitial nephritis: has been reported alone and in association with viral.

Skin and Subcutaneous Tissue Disorders

Photosensitivity reaction

7 DRUG INTERACTIONS

See Table 15. Drug Interactions. SURVEINTE is summarized in this section.

Uridine 5'-diphosphate-glyoxylate (UDG) have been identified as the enzymes responsible for metabolism of lamotrigine. Drugs that induce or inhibit glucuronidation may, therefore, affect the apparent clearance of lamotrigine. Strong or moderate inducers of the cytochrome P450 3A4 (CYP3A4) enzyme, which are also known to induce UDG, may also enhance the metabolism of lamotrigine.

Those drugs that have been demonstrated to have a clinically significant impact on lamotrigine metabolism are outlined in Table 13. Specific dosing guidance for these drugs is provided in the Dosage and Administration section, and for women taking estrogen-containing products, including oral contraceptives, in the Warnings and Precautions section (See Dosage and Administration (2.1), Warnings and Precautions (5.9)).

Additional details of these drug interaction studies are provided in the Clinical Pharmacology section (See Clinical Pharmacology (12.3)).

Concomitant Drug	Effect on Concentration of Lamotrigine or Concomitant Drug	Clinical Comment
Estrogen-containing oral contraceptive preparations containing 30 mcg ethinylloestradiol and 150 mcg levonorgestrel	↓ lamotrigine ↓ levonorgestrel	Decreased lamotrigine concentrations approximately 50%. Decrease in levonorgestrel compared by 19%.
Carbamazepine and carbamazepine epoxide	↓ lamotrigine ? carbamazepine epoxide	Addition of carbamazepine decreases lamotrigine concentration approximately 40%. May increase carbamazepine epoxide levels.
Lopinavir/ritonavir	↓ lamotrigine	Decreased lamotrigine concentration approximately 50%.
Azaxinavir/ritonavir	↓ lamotrigine	Decreased lamotrigine AUC approximately 32%.
Phenobarbital/phenytoin	↓ lamotrigine	Decreased lamotrigine concentration approximately 40%.
Phenytoin	↓ lamotrigine	Decreased lamotrigine concentration approximately 40%.
Ritampin	↓ lamotrigine	Decreased lamotrigine AUC approximately 40%.
Valproate	↓ lamotrigine ↑ valproate	Increase lamotrigine concentrations slightly more than 2-fold. There are conflicting study results regarding effect of lamotrigine on valproate concentrations. In a mean 20% decrease in valproate concentrations in healthy volunteers, 2) no change in valproate concentrations in controlled clinical trials in patients with epilepsy.

↓ Decreased (induces lamotrigine glucuronidation)
↑ Increased (inhibits lamotrigine glucuronidation)
? Conflicting data.

Effect of SURVEINTE on Organic Cations, Transporter 2 Substrates

Lamotrigine is an inhibitor of renal tubular secretion of organic cationic transporter 2 (OCT2) proteins (See Clinical Pharmacology (12.3)). This may result in increased plasma levels of certain drugs that are substantially excreted via these proteins. The combination of SURVEINTE with OCT2 substrates with a narrow therapeutic index (e.g., digoxin) is not recommended.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to AEDs, including SURVEINTE, during pregnancy. Encourage women who are taking SURVEINTE during pregnancy to enroll in the North American Pregnancy Registry (NAEP) Pregnancy Registry by calling 1-888-233-2334 or visiting <http://www.adapregnancyregistry.org>.

Summary

Data from several prospective pregnancy registries and epidemiological studies of pregnant women have not detected an increased frequency of major congenital malformations or a consistent pattern of malformations among women exposed to lamotrigine compared with the general population (see Data). The majority of SURVEINTE pregnancy exposure data are from women with epilepsy. In animal studies, administration of lamotrigine during pregnancy resulted in developmental toxicity (increased mortality, decreased body weight, increased structural variation, neurobehavioral abnormalities) at doses lower than those administered clinically.

Lamotrigine decreased fetal cleft rates in rats, an effect known to be associated with adverse pregnancy outcomes in animals and humans (see Data).

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Clinical Developmental Maternal and/or Embryonal Risk

Epilepsy, with or without exposure to antiepileptic drugs, has been associated with several adverse outcomes during pregnancy, including preterm, low birth weight, placental abruption, placental infarction, poor fetal growth, and perinatal loss. Maternal toxicity was observed in animals, including maternal mortality. The risk of maternal or fetal loss may be greatest for patients with untreated or poorly controlled convulsive seizures.

Women with epilepsy who become pregnant should not abruptly discontinue antiepileptic drugs, and SURVEINTE, due to the risk of status epilepticus or severe seizures, which may be life-threatening (See Warnings and Precautions (5.6)).

Dose Adjustments During Pregnancy and the Postpartum Period

Although with other AEDs, physiological changes during pregnancy may affect lamotrigine concentrations and/or SURVEINTE, due to there have been reports of decreased lamotrigine concentrations during pregnancy and restoration of pre-pregnancy concentrations after delivery. Dose adjustments may be necessary to maintain clinical response.

Data

Human Data: Data from several international pregnancy registries have not shown an increased risk for malformations overall. The International Lamotrigine Pregnancy Registry reported major congenital malformations in 2.2% (95% CI: 1.4, 3.7) versus 1.5% (95% CI: 1.0, 2.0) exposed to lamotrigine monotherapy in the first trimester of pregnancy. The NAEP Pregnancy Registry reported major congenital malformations among 2.9% of 1,562 lamotrigine pregnancies in the first trimester. EUROAP, a large international pregnancy registry focused outside of North America, reported major birth defects in 2.9% (95% CI: 2.3%, 3.7%) of 2,514 exposures to lamotrigine monotherapy in the first trimester. The frequency of major congenital malformations is similar to estimates from the general population.

The NAEP Pregnancy Registry observed an increased risk of isolated oral clefts: among 2,200 infants exposed to lamotrigine early in pregnancy, the risk of oral clefts was 3.2 per 1,000 (95% CI: 1.4, 6.3), a 3-fold increase over the background rate of 1.0 per 1,000. This finding has not been observed in other large international pregnancy registries. Furthermore, a case-control study based on 21 congenital anomaly registries covering over 10 million births in Europe reported an adjusted odds ratio for isolated oral clefts with lamotrigine exposure of 1.45 (95% CI: 0.8, 2.6).

Several meta-analyses have not reported an increased risk of major congenital malformations following lamotrigine monotherapy in pregnancy compared with healthy and disease-matched controls. No patterns of specific malformation types were observed.

The same meta-analyses evaluated the risk of additional maternal and infant outcomes including fetal death, stillbirth, preterm birth, small for gestational age, and neurodevelopmental delay. Although there are no data suggesting an increased risk of these outcomes with lamotrigine monotherapy exposure, differences in outcome definition, ascertainment methods, and comparator groups limit the conclusions that can be drawn.

Animal Data: When lamotrigine was administered to pregnant mice, rats, or rabbits during the period of organogenesis (oral doses of up to 125, 25, and 50 mg/kg, respectively), reduced fetal body weight and increased incidence of fetal skeletal variations were seen in mice and rats at doses that were also maternally toxic. The effect doses for embryofetal developmental toxicity in mice, rats, and rabbits (75, 625, and 50 mg/kg, respectively) are similar to (mice and rabbits) or less than (the human dose of 400 mg/day on a body surface area (m²) basis).

In a study in which pregnant rats were administered lamotrigine (oral doses of 5, 10, or 20 mg/kg) during the latter part of gestation and throughout lactation, increased offspring mortality (including stillbirths) was seen at all doses. The lowest effect dose for pre- and post-natal losses tested during lactation in rats is less than the human dose of 400 mg/day on a m² basis. Maternal toxicity was observed at the 2 highest doses tested.

When administered in pregnant rats, lamotrigine decreased fetal body concentrations at doses greater than or equal to 5 mg/kg/day, which is less than the human dose of 400 mg/day on a m² basis.

8.2 Lactation

Risk Summary

Lamotrigine is present in milk from lactating women taking SURVEINTE (see Data). Neonates and young infants are at risk for high plasma levels because maternal serum and milk levels can rise to high levels postpartum if lamotrigine has been increased during pregnancy and is not reduced after delivery. Glucuronidation is required for drug clearance. Glucuronidation capacity is immature in the infant and this may also contribute to the level of lamotrigine exposure. Events including rash, anorexia, drowsiness, poor sucking, and poor weight gain (resulting in hospitalization in some cases) have been reported in human milk-fed infants by mothers using lamotrigine, whether or not these events were caused by lamotrigine is unknown. No data are available on the effects of the drug on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for SURVEINTE and any potential adverse effects on the breastfed infant from SURVEINTE or from the underlying maternal condition.

Clinical Considerations

Human milk-fed infants should be closely monitored for adverse effects resulting from lamotrigine. Measurement of infant serum levels should be performed to rule out toxicity if concerns arise. Human milk-testing should be discontinued in infants with lamotrigine toxicity.

Data

Data from multiple adult studies indicate that lamotrigine plasma levels in nursing infants have been reported to be as high as 50% of maternal plasma concentrations.

8.4 Pediatric Use

Epilepsy

SURVEINTE is indicated as adjunctive therapy in patients aged 2 years and older for partial-onset seizures, the generalized seizures of Lennox-Gastaut syndrome, and PGTs.

Safety and efficacy of SURVEINTE used as adjunctive treatment for partial-onset seizures were not demonstrated in a small, randomized, double-blind, placebo-controlled withdrawal trial in very young pediatric patients (aged 1 to 24 months). SURVEINTE was associated with an increased risk for infectious adverse reactions (SURVEINTE 2%, placebo 2%), and respiratory adverse reactions (SURVEINTE 8%, placebo 5%). Infectious adverse reactions included bronchitis, bronchiolitis, ear infection, eye infection, otitis externa, pharyngitis, urinary tract infection, and viral infection. Respiratory adverse reactions included nasal congestion, cough, and croup.

Bleeding Disorder

Safety and efficacy of SURVEINTE for the maintenance treatment of bipolar disorder were not established in a double-blind, randomized withdrawal, placebo-controlled trial that evaluated 201 pediatric patients aged 10 to 17 years with a current manic/mixed episode, depressed or mixed mood episode as defined by DSM-IV-TR. In the randomized phase of the trial, adverse reactions that occurred at least 5% of patients taking SURVEINTE (n = 47) and were twice as common compared with patients taking placebo (n = 89) were: viral infections (SURVEINTE 8%, placebo 2%), oropharyngeal pain (SURVEINTE 8%, placebo 2%), vomiting (SURVEINTE 8%, placebo 2%), contact dermatitis (SURVEINTE 5%, placebo 2%), upper abdominal pain (SURVEINTE 5%, placebo 1%), and suicidal ideation (SURVEINTE 5%, placebo 0%).

Juvenile Animal Data

In a juvenile animal study in which lamotrigine (oral doses of 0.5, 15, or 20 mg/kg) was administered to young rats from postnatal day 7 to 62, decreased viability and growth were seen at the highest dose tested and long-term neurobehavioral abnormalities developed including hyperactivity, increased reactivity, and learning effects in animals tested as adults) were observed at the 2 highest doses tested. The no-effect dose for adverse developmental effects in juvenile animals was less than the human dose of 400 mg/day on a m² basis.

5.5 Geriatric Use

Clinical trials of SURVEINTE for epilepsy and bipolar disorder did not include sufficient numbers of patients aged 65 years and older to determine whether they respond differently from younger patients or require a different safety profile than that of younger patients. Caution should be used when prescribing SURVEINTE to geriatric patients, especially during the low end of the dosing range, reflecting the greater frequency of decreased kidney, liver, or cardiac function and of concomitant disease or other drug therapy.

5.6 Hepatic Impairment

Exposure in patients with hepatic impairment is limited. Based on a clinical pharmacology study in 24 subjects with mild, moderate, and severe liver impairment, the following general recommendations can be made (See Clinical Pharmacology (12.3)). No dosage adjustment is needed in patients with mild liver impairment. Initial, escalation, and maintenance doses should generally be reduced by approximately 50% in patients with moderate and severe liver impairment without cirrhosis and 50% in patients with severe liver impairment without cirrhosis. Escalation and maintenance doses may be adjusted according to clinical response (See Dosage and Administration (2.1)).

5.7 Renal Impairment

SURVEINTE is metabolized mainly by glucuronic acid conjugation, with the majority of the metabolites being recovered in the urine. In a small study comparing a single dose of lamotrigine in subjects with varying degrees of renal impairment with healthy volunteers, the plasma half-life of lamotrigine was approximately twice as long as the subjects with chronic renal failure (See Clinical Pharmacology (12.3)).

10 OVERDOSAGE

10.1 Human Overdose Experience


Overdoses involving quantities up to 15 g have been reported for SURVEINTE, some of which have been fatal. Overdose has resulted in ataxia, nystagmus, seizures (including tonic-clonic seizures), decreased level of consciousness, coma, and intracranial conduction delay.

10.2 Management of Overdose

There are no specific antidotes for lamotrigine. Following a suspected overdose, hospitalization of the patient is advised. General supportive care is indicated, including frequent monitoring of vital signs and close observation of the patient. If indicated, emesis should be induced; usual precautions should be taken to protect the airway. It should be kept in mind that immediate-release lamotrigine is rapidly absorbed (See Clinical Pharmacology (12.3)). If uncertain whether lamotrigine is an effective means of removing lamotrigine from the blood, in 6 renal failure patients, about 20% of the amount of lamotrigine in the body was removed by hemodialysis during a 4-hour session. A Poison Control Center should be contacted for information on the management of overdose of SURVEINTE.

11 DESCRIPTION

Lamotrigine, USP, an AED of the phenylhydantoin class, is chemically unrelated to existing AEDs. Lamotrigine's chemical name is 3-(5-dimethyl-6-(2,3-dichlorophenyl)-dehydro-1H, its molecular formula is C₁₄H₁₁Cl₂N₃, and its molecular weight is 266.09. Lamotrigine, USP is a white to pale cream-colored powder and has a pK_a of 5.7. Lamotrigine, USP is very slightly soluble in water (0.17 mg/mL at 25°C) and slightly soluble in 0.1 M HCl (4.1 mg/mL at 25°C). The structural formula is:



SURVEINTE (lamotrigine) tablets, USP are supplied for oral administration as 25-mg (white to off white), 100-mg (white to off white), 150-mg (white to off white), and 200-mg (white to off white) tablets. Each tablet contains the labeled amount of lamotrigine, USP and the following inactive ingredients: lactose monohydrate, magnesium stearate, microcrystalline cellulose, povidone, and sodium starch glycolate.

Meets USP Dissolution Test 1

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The precise mechanism(s) by which lamotrigine exerts its anticonvulsant action are unknown. In animal models designed to detect anticonvulsant activity, lamotrigine was effective in preventing seizure spread in the maximum electroshock (MES) and pentylenetetrazol (scotized) tests, and prevented seizures in the visually and electrically evoked after-discharge (EAD) tests for antiepileptic activity. Lamotrigine also displayed embryonic protection in the kindling model of rats both during kindling development and in the fully kindled state. The relevance to these models to human epilepsy, however, is not known.

One proposed mechanism of action of lamotrigine, the relevance of which remains to be established in humans, involves an effect on sodium channels. In vitro pharmacological studies suggest that lamotrigine inhibits voltage-sensitive sodium channels, thereby stabilizing neuronal membranes and consequently decreasing neuronal excitability. Excitatory amino acids (e.g., glutamate and aspartate).

Effect of Lamotrigine on N-Methyl-D-Aspartate Receptor-Mediated Activity

Lamotrigine did not inhibit N-methyl-D-aspartate (NMDA)-induced depolarizations in rat cortical slices or NMDA-induced cyclic GMP formation in rat cerebellar slices. Lamotrigine did not inhibit NMDA-induced depolarizations in rat cortical slices or NMDA-induced cyclic GMP formation in rat cerebellar slices. Lamotrigine did not inhibit NMDA-induced depolarizations in rat cortical slices or NMDA-induced cyclic GMP formation in rat cerebellar slices.

12.2 Pharmacokinetics

Folate Metabolism

In vitro, lamotrigine inhibited dihydrofolate reductase, the enzyme that catalyzes the reduction of dihydrofolate to tetrahydrofolate. Inhibition of this enzyme may interfere with the biosynthesis of nucleic acids and other biologically important molecules. However, in clinical studies, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Cardiac Electrophysiology

Effect of Lamotrigine: In vitro studies show that lamotrigine exhibits Class IB antiarrhythmic activity at therapeutically relevant concentrations. Lamotrigine is an inhibitor of the hERG potassium current, but it does not block the hERG current at clinically relevant concentrations. Lamotrigine is an inhibitor of the hERG potassium current, but it does not block the hERG current at clinically relevant concentrations.

In vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabolite: In vitro, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. The in vivo electrophysiological effects of this metabolite have not been studied. Similar cardiovascular effects from this metabolite are not expected to be observed in humans. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism. In patients with mild to moderate renal impairment, lamotrigine had no effect on folate metabolism.

Effect of Lamotrigine Metabol